

(3 Hours)

[Total Marks : 80]

**Note:**

- 1) Question No.1 is compulsory
- 2) Attempt any three out of remaining five questions
- 3) Figures to the right indicate full marks

**Q1.**

- a) If  $\sin(\theta + i\phi) = \tan\alpha + i\sec\alpha$ , then show that  $\cos 2\theta \cdot \cosh 2\phi = 3$  [5]
- b) If  $u = \log(\tan x + \tanh y)$ , then show that  $\sin 2x \frac{\partial u}{\partial x} + \sin 2y \frac{\partial u}{\partial y} = 2$  [5]
- c) Express the matrix  $A = \begin{bmatrix} 0 & 5 & -3 \\ 1 & 1 & 1 \\ 4 & 5 & 9 \end{bmatrix}$  as the sum of a symmetric and skew symmetric matrix. [5]
- d) Expand  $\sqrt{1 + \sin x}$  in ascending powers of  $x$  upto  $x^4$  term. [5]

**Q2.**

- a) Find non-singular matrices P and Q such that PAQ is in normal form where, [6]

$$A = \begin{bmatrix} 4 & 3 & 1 & 6 \\ 2 & 4 & 2 & 2 \\ 12 & 14 & 5 & 16 \end{bmatrix}. \text{ Also find the rank of A.}$$

- b) If  $z = f(x, y)$  and  $x = u \cosh v$ ,  $y = u \sinh v$ ; prove that [6]

$$\left(\frac{\partial z}{\partial x}\right)^2 - \left(\frac{\partial z}{\partial y}\right)^2 = \left(\frac{\partial z}{\partial u}\right)^2 - \frac{1}{u^2} \left(\frac{\partial z}{\partial v}\right)^2$$

- c) Prove that  $\text{Log} \left[ \frac{(a-b)+i(a+b)}{(a+b)+i(a-b)} \right] = i(2n\pi + \tan^{-1} \frac{2ab}{a^2-b^2})$ . Hence evaluate  $\text{Log} \left( \frac{1+5i}{5+i} \right)$  [6]

Q3.

- a) If  $\alpha$  and  $\beta$  are the roots of the equation  $z^2 \sin^2 \theta - z \sin 2\theta + 1 = 0$ , then prove that  
 $\alpha^n + \beta^n = 2 \cos n\theta \operatorname{cosec}^{2n} \theta$  and  $\alpha^n \beta^n = \operatorname{cosec}^{2n} \theta$  [6]
- b) Solve the following equations by Gauss-Seidal Method ; [6]  
 $15x + 2y + z = 18$  ,  $2x + 20y - 3z = 19$ ,  $3x - 6y + 25z = 22$ ,  
 Take three iterations.

- c) Prove that if  $z$  is a homogeneous function of two variables  $x$  and  $y$  of degree  $n$ , then  
 $x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = n(n-1)z$ . Hence find the value of  $x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2}$   
 at  $x = 1, y = 1$  when  $z = x^6 \tan^{-1} \left( \frac{x^2 + y^2}{x^2 + xy} \right) + \frac{x^4 + y^4}{x^2 y^2}$  [8]

Q4.

- a) If  $\tan(\alpha + i\beta) = \cos \theta + i \sin \theta$  then prove that  $\alpha = \frac{n\pi}{2} + \frac{\pi}{4}$ ,  $\beta = \frac{1}{2} \log \left( \frac{\pi}{4} + \frac{\theta}{2} \right)$  [6]
- b) Expand  $x^5 + x^3 - x^2 + x - 1$  in powers of  $(x - 1)$  and hence find the value of [6]  
 1)  $f\left(\frac{9}{10}\right)$   
 2)  $f(1.01)$
- c) For what values of  $\lambda$  and  $\mu$ , the equations, [8]  
 $x + y + z = 6$  ;  $x + 2y + 3z = 10$  ;  $x + 2y + \lambda z = \mu$   
 1) have a unique solution  
 2) have infinite solution

Find the solution in each case for a possible value of  $\mu$  and  $\lambda$ .

**Q5.**

a) Find the nth derivative of  $y = \frac{1}{x^2 + a^2}$  [6]

b) Discuss the maxima and minima of  $x^3 + xy^2 - 12x^2 - 2y^2 + 21x + 16$  [6]

c) Prove that if A and B are two unitary matrices then AB is also unitary. Verify the result when

$$A = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1+i \\ 1-i & -1 \end{bmatrix} \text{ and } B = \begin{bmatrix} \frac{1+i}{2} & \frac{-1+i}{2} \\ \frac{1+i}{2} & \frac{1-i}{2} \end{bmatrix} \quad [8]$$

**Q6.**

a) If  $x = \cosh\left(\frac{1}{m} \log y\right)$ , prove that [6]

$$(x^2 - 1)y_{n+2} + (2n + 1)x y_{n+1} + (n^2 - m^2) y_n = 0$$

b) Find a root of the equation  $x e^x = \cos x$  using the Regular Falsi Method correct to three decimal places. [6]

c) 1) Expand  $\sin^4 \theta \cos^2 \theta$  in a series of multiples of  $\theta$ . [4]

2) If one root of  $x^4 - 6x^3 + 18x^2 - 24x + 16 = 0$  is  $(1+i)$ ; find the other roots. [4]